

Important Safety Information for Gastroenterologists and Hepatologists About Potential Risks of Gastrointestinal Perforation and Transaminase Elevations With ACTEMRA®

ACTEMRA® (tocilizumab) is an interleukin-6 (IL-6) receptor inhibitor that has been approved by the Food and Drug Administration (FDA) for two indications:

- Adult patients with moderately to severely active rheumatoid arthritis (RA) who have had inadequate response to one or more TNF antagonist therapies with a recommended ACTEMRA® dosing interval of every 4 weeks.
- Children 2 years of age and older with active *Systemic Juvenile Idiopathic Arthritis (SJIA)* with a recommended ACTEMRA® dosing interval of every 2 weeks.

The safety and efficacy of ACTEMRA® for conditions other than RA and SJIA have not yet been established.

Gastroenterologists and **hepatologists** should be aware of important safety information regarding ACTEMRA.

Gastrointestinal perforations: Gastrointestinal (GI) perforations have been reported in Phase 3 clinical trials, primarily as complications of diverticulitis, including generalized purulent peritonitis, lower GI perforation, fistula and abscess. Most patients who developed GI perforations were taking concomitant nonsteroidal anti-inflammatory medications (NSAIDs), corticosteroids or methotrexate. ACTEMRA should be used with caution in patients who may be at increased risk for GI perforation. Patients presenting with new-onset abdominal symptoms should be evaluated promptly for early identification of GI perforation.

Transaminase elevations: Treatment with ACTEMRA was associated with a higher incidence of transaminase elevations (ALT, AST) in Phase 3 clinical trials. These elevations did not result in apparent permanent or clinically evident hepatic injury with modification of the treatment regimen, which resulted in a decrease or normalization of liver enzymes. Patients receiving ACTEMRA should be monitored for elevated transaminase levels and dose modifications may be necessary. When clinically indicated, other liver function tests, such as bilirubin, should be considered. Please see full Prescribing Information for more information.

Reporting Adverse Events

It is important that you report any serious gastrointestinal adverse events, including GI perforation, hepatic disease or hepatic impairment, that occur in a patient being treated with ACTEMRA, even if you do not think there is a causal relationship. The information that you, as a gastroenterologist or hepatologist, provide about these events may inform therapy and monitoring decisions for future patients.

Reporting is easy and maintains patient confidentiality. Your patient's name or contact information is not needed. *HIPAA does not apply to this adverse event reporting.* You can report your cases to Genentech or directly to the FDA:

- Genentech at 1-800-ACTEMRA (1-800-228-3672)
- MedWatch (FDA safety information and adverse event reporting program) at 1-800-332-1088 or online at www.fda.gov/medwatch/report.htm

Please visit www.ACTEMRA.com for full Prescribing Information, including Boxed Warning, and Medication Guide.